



RACC-TRIAL postoperative visit – 2 years

Institution/Hospital _____ City _____ Country _____

Date of visit: _____

Patient status:

- Alive Dead Not known
- No evidence of disease Evidence of disease Relapse has occurred before visit

CTCAE 3.0 (Lymphatic side effects according to the Common Terminology Criteria (CTC) version 3.0)

Please evaluate the current status of the patient and put X in one of the boxes

Adverse event	None	Grade 1	Grade 2	Grade 3	Grade 4
Lower extremity					
Truncal/genital					
lymphocele					

Adverse event	Grade 1	Grade 2	Grade 3	Grade 4
Lower extremity lymphedema	5%–10% inter-limb discrepancy in volume or circumference at point of greatest visible difference; pitting edema	10–30% inter-limb discrepancy in volume or circumference at point of greatest visible difference	>30% inter-limb discrepancy in volume; lymphorrhea, interfering with activities of daily life	Progression to malignancy (i.e. lymphangiosarcoma), amputation indicated, disabling
Truncal/genital lymphedema	Swelling or obscuration of anatomic architecture on close inspection, pitting edema	Readily apparent obscuration of anatomic architecture; obliteration of skin folds	Lymphorrhea; interfering with activities of daily life; gross deviation from normal anatomical contour	Progression to malignancy (i.e. lymphangiosarcoma); disabling
Lymphocele	Asymptomatic, clinical or radiographic findings only	Symptomatic; medical intervention indicated	Symptomatic and interventional radiology or operative intervention indicated	–

Quality of life questionnaire:

- yes no the patient has filled in the Quality of Life questionnaire

Date of completion _____